BNS-042

No. of Printed Pages: 16

CERTIFICATE IN COMMUNITY HEALTH FOR NURSES (BPCCHN)

Term End Examination,

December, 2021

BNS-042: PRIMARY HEALTH CARE IN COMMON CONDITIONS

Time: 2 Hours Maximum Marks: 70

GENERAL INSTRUCTIONS

- 1. All questions are **compulsory**. Each question carries 1 mark.
- 2. No cell phones, calculators, books, slide-rules, notebooks or written notes, etc. will be allowed inside the examination hall.
- 3. You should follow the instructions given by the Centre Superintendent and by the Invigilator at the examination venue. If you violate the instructions, you will be disqualified.
- 4. Any candidate found copying or receiving or giving assistance in the examination will be disqualified.
- 5. The Question Paper and the OMR Response Sheet (Answer Sheet) would be supplied to you by the Invigilators. After the examination is over, you should hand over the OMR Response Sheet to the Invigilator before leaving the examination hall. Any candidate who does not return the OMR Response Sheet will be disqualified and the University may take further action against him/her.
- 6. All rough work is to be done on the question paper itself and not on any other paper. Scrap paper is not permitted. For arriving at answers you may work in the margins, make some markings or underline in the test booklet itself.
- 7. The University reserves the right to cancel the result of any candidate who impersonates or uses/adopts other malpractices or uses any unfair means. The University may also follow a procedure to verify the validity of scores of all examinees uniformly. If there is substantial indication that your performance is not genuine, the University may cancel your result.

How to fill up the information on the OMR Response Sheet (Examination Answer Sheet)

- 1. Write your complete Enrolment No. in 9 digits. This should correspond to the enrolment number indicated by you on the OMR Response Sheet. Also write your correct name, address with pin code in the space provided. Put your signatures on the OMR Response Sheet with date. Ensure that the Invigilator in your examination hall also puts his signatures with date on the OMR Response Sheet at the space provided.
- 2. On the OMR Response Sheet student's particulars are to be filled in by blue/black ball pen also. Use blue/black ball pen for writing the Enrolment No. and Examination Centre Code as well as for blackening the circle bearing the correct answer number against the serial number of the question.
- 3. Do not make any stray remarks on this sheet.
- 4. Write correct information in numerical digits in Enrolment No. and Examination Centre Code Columns. The corresponding circle should be dark enough and should be filled in completely.
- 5. Each question is followed by four probable answers which are numbered (1), (2), (3) and (4). You should select and show only one answer to each question considered by you as the most appropriate or the correct answer. Select the most appropriate answer. Then by using blue/black ball pen, blacken the circle bearing the correct answer number against the serial number of the question. If you find that answer to any question is none of the four alternatives given under the question, you should darken the circle with '0'.
- 6. No credit will be given if more than one answer is given for one question. Therefore, you should select the most appropriate answer.
- 7. You should not spend too much time on one question. If you find any particular question difficult, leave it and go to the next. If you have time left after answering all the questions, you may go back to the unanswered question.
- 8. There is no negative marking for wrong answers.

BNS-042 (2)

1.	Jaur	ndice with right hypochondrial p	ain n	nay be related to:
	(1)	Hepatitis-A infection	(2)	Gall bladder stone
	(3)	Hepatitis-B infection	(4)	Hepatitis-C infection
2.	The	sign and symptoms of upper gas	stroin	testinal bleeding of the following except:
	(1)	Hypertension	(2)	Hematemesis
	(3)	Coffee ground gastric aspirate	(4)	Tachycardia
3.	All c	of the following are causes of abo	lomin	al distension except :
	(1)	Intestinal perforation	(2)	Peritonitis
	(3)	Diabetes mellitus	(4)	Cirrhosis of liver
4.	A pa	atient with dyspepsia would be a	dvise	d all the following measures except :
	(1)	Avoid excess tea, coffee	(2)	Stop alcohol and smoking
	(3)	Avoid spicy food	(4)	Follow meal at any time
5.	Diet	ary measures for upper respi	rator	y infections include all of the following
	exce	pt:		
	(1)	Vitamin C	(2)	Refined carbohydrates
	(3)	Chicken soup	(4)	Sprouts
6.	A pa	atient with sinusitis should avoid	d:	
	(1)	Decongestant spray	(2)	Pain killer
	(3)	Antibiotics	(4)	Steam inhalation
BNS-	042		(3)	

7.	Acut	te chest pain is common sympton	n in a	all of the following conditions <i>except</i> :
	(1)	DVT	(2)	Pulmonary embolism
	(3)	Heart diseases	(4)	Anxiety disorders
8.	One	of the natural home remedies for	r pat	ient with bronchial asthma includes :
	(1)	Start I/V infusion		
	(2)	Rubbing of mustard oil-campho	or mix	xed on back
	(3)	Start humidified oxygen therap	у	
	(4)	Give cold water to drink		
9.	All o	of the following conditions are re	lated	to urinary tract infections $except$:
	(1)	Cystitis	(2)	Pyelonephritis
	(3)	Vagitinitis	(4)	Prostatis
10.	One	of the first aid measure for conj	uncti	vitis includes :
	(1)	Do not bandage the eye withou	t kno	wing cause
	(2)	Torch light examination		
	(3)	Intraocular pressure measuren	nent	
	(4)	Fundus examination		
11.	1. The shock which occurs immediately after injury is :			
	(1)	Acute	(2)	Primary shock
	(3)	Chronic	(4)	Secondary shock

(4)

12.	One	e of the following is true about haemo	rrhagic stroke :
	(1)	Similar to heart attack	
	(2)	Clot forms in the blood vessels	
	(3)	Rupture of blood vessels in the brai	n
	(4)	Fatty deposits clogs the blood vesse	ls
13.	A sc	cheme launched to motivate women t	o deliver at health facility is :
	(1)	Janani Suraksha Karyakram (JSK)	
	(2)	Janani Shishu Suraksha Karyakra	m
	(3)	Navjat Shishu Janani Suraksha Ka	ryakram
	(4)	Janani Suraksha Yojna (JSY)	
14.	New	w initiatives to strengthen routine i	mmunization involves all of the following
	exce	ept:	
	(1)	Auto Disabled (AD) syringes	
	(2)	Maintaining cold chain only on the	day of immunization
	(3)	Mobilization of children by ASHA	
	(4)	Alternate vaccine delivery from PH	C to subcenters
15.	One	e of the following related to colou	r coded dashboard under HMIS (Health
	Mar	nagement Information System) is not	true:
	(1)	Very good (light green) (2)	Good (green)
	(3)	Promising (yellow) (4)	Very poor performing (red)
BNS-	-042	(5)	

	(1)	Deciding to seek care			
	(1)	Deciding to seek care			
(2) Identifying and reaching health facility					
(3) Receiving appropriate treatment					
	(4)	Accepting inadequate care from	n heal	th facility	
17.	A p	regnant woman attending the	e hea	alth facility with vomiting during first	
	trim	aster requires one of the following	ng ac	tion:	
	(1)	to eat small frequent meals.			
	(2)	start I/V ringer lactate.			
	(3)	refer her to MO for further man	nagen	nent.	
	(4)	advise her to have a hospital de	eliver	y.	
18.	All t	he following are true about TT i	nfecti	ion $except$:	
	(1)	Dosage is 0.5 ml	(2)	Given in gluteal muscles	
	(3)	Deep intramuscular	(4)	Given in the upper arm	
19.	The	daily Recommended Dietary A	llowa	ance (RDA) for calcium in pregnancy and	
	lacta	ation is:			
	(1)	400 mg/day	(2)	600 mg/day	
	(3)	1200 mg/day	(4)	1000 mg/day	
20.	Fetu	as is dependent on which of the	follov	wing maternal hormones in first trimester	
	of pr	regnancy?			
	(1)	Insulin	(2)	Glucagon	
	(3)	Progesterone	(4)	Maternal transplacental thyroid hormone	
BNS-	042		(6)		

16. All of the following delay the seeking of MCH services *except*:

21.	Prot	cocol for management of hypot	hyroidi	$egin{aligned} ext{dism includes all of the following } except: \end{aligned}$			
	(1)	Drug to be taken half an hou	r after	r lunch			
	(2)	Drug of choice is tablet levotl	hyroxin	ine			
	(3)	Drug to be taken in morning-	empty	ty stomach			
	(4)	Drug can be used safely duri	ng preg	egnancy and lactation			
22.	Trea	atment for hypothyroidism s	should	l be discontinued, when level of TSH is			
	with	in :					
	(1)	1-8 ml U/L	(2)	2-9 ml U/L			
	(3)	3-10 ml U/L	(4)	Above 10 ml U/L			
23.	For	treating syphilis during pregn	ancy, n	most suitable drug is:			
	(1)	Amoxicillin	(2)	Penicillin			
	(3)	Cloxacin	(4)	Amikacin			
24.	All t	the following are true about no	ormal la	${ m labour}\ except:$			
	(1)	Between 37 to 42 completed weeks of gestation					
	(2)	Spontaneous onset					
	(3)	Cephalic presentation					
	(4)	Completes within 18 hours or	r more	e			
25.	Mag	gnesium sulphate is the drug o	f choice	ce for treatment of :			
	(1)	Eclampsia	(2)	Antepartum haemorrhage			
	(3)	Post-partum haemorrhage	(4)	Retained placenta			
BNS-	042		(7))			

26.	WH	O recommendation for spacing a	ıfter a	miscarriage or induced abortion is:
	(1)	3 months	(2)	at least 6 months
	(3)	4 months	(4)	at least 5 months
27.	All	the following are causes of secon	dary	$ ext{dysmenorrhoea}\ except:$
	(1)	Uterine fibroid	(2)	Cervical narrowing
	(3)	Nulliparity	(4)	Ovarion-abscess
28.	All	of the following are common	nly p	rescribed medications for premenstrual
	syno	$drome\ except:$		
	(1)	Antiviral	(2)	Diuretics
	(3)	Analgesics	(4)	Antidepressants
29.	Thic	ck, white, cheesy vaginal dischar	ge is	caused by:
	(1)	Gonorrhoea	(2)	Trichomoniasis
	(3)	Bacterial infection	(4)	Yeast infection
30.	All	of the following preventive meas	sures	should be taken for prevention of mastitis
	exce	pt:		
	(1)	Equal breast feed from both br	easts	
	(2)	Affly cold compress to the brea	st	
	(3)	Avoid dehydration by drinking	plent	ty of fluids
	(4)	Wear a supportive bra		

(8)

31.	One of the following is a type of vaginal prolapse:								
	(1)	Cystocele	(2)	Released perineum					
	(3)	Rectocele	(4)	Enterocele					
32.	Othe	er name for ovulation method is	:						
	(1)	Basal body temperature method							
	(2)	Calender method							
	(3)	Cervical mucus method							
	(4)	Standard days method							
33.	An C	Oral Contraceptive Pills (OCPs)	packe	et contains :					
(1) 7 white pills of contraceptive + 21 brown pills				rown pills					
(2) 21 white pills of contraceptive + 7 brown pills (icon)									
	(3) 20 white pills of contraceptive + 6 brown pills								
	(4) 30 white pills of contraceptive + 7 brown pills								
34.	All o	f the following are advantages o	of infe	ectable contraceptives <i>except</i> :					
	(1)	Highly effective	(2)	Reversible					
	(3)	Long acting	(4)	Irreversible					
35.	Und	er Prerna strategy, B. P. L. coup	oles a	re awarded :					
	(1)	₹ 9,000 for boy or ₹ 10,000 for g	girl ch	aild					
	(2)	₹ 10,000 at birth of a boy or ₹ 1	2,000	for girl					
	(3)	₹ 8,000 for boy or ₹ 9,000 for gi	rl chil	ld					
	(4)	₹ 6,000 for body or ₹ 8,000 for g	girl ch	aild					

(9)

36.	All of the following are principles of counselling except :				
	(1) Counsellor has right to take decision				
	(2)	Client has right to choose action	ons		
	(3) Strictly confidential				
	(4) Helps client to identify the problem				
37.	Cor	rect sequence of counselling sess	sion is	3:	
	(1)	Connect-Reassure-Stabilise-Fa	ıcilita	te Coping	
	(2) Reassure-Connect-Stabilise-Facilitate Coping				
	(3)	Stabilise-Reassure-Connect-Fa	acilita	te Coping	
	(4)	Facilitate Coping-Stabilise-Rea	assur	e-Connect	
38.	Rate	e of teenage pregnancy in India	is:		
	(1)	102 per 1000 women	(2)	202 per 1000 women	
	(3)	62 per 1000 women	(4)	152 per 1000 women	
39.	All	of the following components of	of wa	rm chain are important during delivery	
	exce	pt:			
	(1)	Delivery room temperature is	25°C.		
	(2)	Dry the baby with wet cloth.			
	(3)	Cover baby with clean dry clot	h.		
	(4)	Skin to skin contact with moth	er.		
40.	Res	piratory rate of a normal newbo	rn bal	by is:	
	(1)	40 to 60 times per minute	(2)	30 to 50 times per minute	
	(3)	20–40 times per minute	(4)	25–40 times per minute	

(10)

41.	Tak	ing out of curdled milk soon afte	er feed	d is known as :
	(1)	Aspiration	(2)	Inhalation
	(3)	Regurgitation	(4)	Exhalation
42.	All	of the following diseases are cau	sed by	y Staphylococcol and Streptococcol <i>except</i> :
	(1)	Impetigo	(2)	Cellulitis
	(3)	Folliculitis	(4)	Vertigo
43.	Asse	essment of sick young infant inv	olves	all of the following skills $except$:
	(1)	Classifying	(2)	Asking
	(3)	Looking at	(4)	Listening
44.	All	of the following points should	be k	kept in mind while giving intramuscular
	anti	$biotics\ except:$		
	(1)	Explain to the mother why the	drug	is given.
	(2)	Determine the dose of antibiot	ic.	
	(3)	Administer the drug with clear	need	dle and syringe.
	(4)	Administer the drug with steri	le aut	to-disposable syringe.
45.	A co	ondition in which placenta is in	nplan	ted partially or completely over the lower
	uter	rine segment is called:		
	(1)	Placenta accreta	(2)	Placenta praevia
	(3)	Abruptio placenta	(4)	Placenta circumvalleta
46.	Con	nmon drug to treat both pneumo	nia a	nd dysentery is:
	(1)	Cotrimaxazole	(2)	Gentamicin
	(3)	Amoxicillin	(4)	Penicillin
BNS-	042		(11)	

47.	All	of the following are birth defects	excep	t:
	(1)	Neural tube defects	(2)	Down's syndrome
	(3)	Rickets	(4)	Congenital heart defects
48.	All	of the following are disabilities a	ımong	children except :
	(1)	Vision impairment	(2)	Hearing impairment
	(3)	Language impairment	(4)	Convulsion
49.	Cou	ntry having largest coverage of	unive	rsal immunization programme is :
	(1)	China	(2)	India
	(3)	Bangladesh	(4)	Sri Lanka
50.	Gov	ernment of India launched Miss	ion In	adradhanush to achieve :
	(1)	To reduce maternal mortality	(2)	To reduce infant mortality
	(3)	Full immunization coverage	(4)	To reduced malnutrition in children
51.	One	of the following vaccines is sens	sitive	to freezing :
	(1)	BCG	(2)	Measles
	(3)	Hepatitis-B	(4)	OPV
52.	All	of the following vaccines can	cause	anaphylactic shock after administration
	exce	pt:		
	(1)	Hepatitis-B	(2)	Measles
	(3)	Tetanus	(4)	BCG
53.	All	of the following infections result	from	unsafe injection practices <i>except</i> :
	(1)	Hepatitis-B	(2)	HIV/AIDS
	(3)	Hepatitis-A	(4)	Hepatitis-C
BNS-	042		(12)	

54.	To p	orepare hypochlorite solution at	home	use one of the following:					
	(1)	mix 2 table spoons of bleaching	g pow	der with coil					
	(2)	(2) dissolve 10-15 gm of bleaching powder in one litre of water.							
	(3)	(3) one teaspoon of bleaching powder with 1 litre of water.							
	(4)	freshly diluted solution of salt	and v	vater.					
55.	All	of the following needs to be	cleare	ed from an injured person's mouth after					
	mee	eting safety need except:							
	(1)	any blood	(2)	vomit					
	(3)	any foreign body	(4)	pus					
56.	The	size of the needle required to as	perat	te the fluid in haemothorax is:					
	(1)	14 or 16 gauze needle	(2)	16 or 18 gauze needle					
	(3)	18 or 20 gauze needle	(4)	24 or 26 gauze needle					
57.	One	of the following disease is	1009	% fatal, 100% vaccine preventable and					
	incu	rable :							
	(1)	Scabies	(2)	Scoliosis					
	(3)	Rabies	(4)	Pneumoconiosis					
58.	A pa	atient with severe epigastric pai	n rad	iating to back is :					
	(1)	Acute pancreatitis	(2)	Peptic ulcer					
	(3)	Duodenal ulcer	(4)	Chronic pancreatitis					
59.	The	choice of drug for treatment of t	racho	oma is :					
	(1)	Doxycycline ointment	(2)	Tetracycline eye ointment					
	(3)	Tetracycline oral capsules	(4)	Doxycycline oral capsules					
BNS-	042		(13)						

60.	. In which of the following conditions there is protrusion of abdominal contents			
	thro	ugh a sac outside the abdominal	l wall	?
	(1)	Varicocele	(2)	Hydrocele
	(3)	Hernia	(4)	Lymphedema
61.	Dysi	phagia is known as :		
	(1)	Swelling of glands	(2)	Difficulty in swallowing
	(3)	Difficulty in eating	(4)	Difficulty in drinking
62.	All o	of the following are true about at	rophi	c vaginitis except :
	(1)	Occur in post-menopausal wom	ien	
	(2)	Thinning and drying of vaginal	wall	
	(3)	Dyspareunia		
	(4)	Painless intercourse		
63.	Tali	pes equinus is also commonly kr	nown	as:
	(1)	Club foot	(2)	Exomphalos major
	(3)	Exomphalos minor	(4)	Gastroschisis
64.	Palli	iative care involves :		
	(1)	Superspeciality approach	(2)	Multi-disciplinary team approach
	(3)	Specific team approach	(4)	Speciality care approach
65.	Whi	ch one of the following diet redu	ces th	ne risk of cancer?
	(1)	Animal fats	(2)	Salted foods
	(3)	Rich in fibre	(4)	High sugar foods
BNS-	042		(14)	

66.	Drug used as a prophylaxis of post-operative nausea:			
	(1)	Domperidone	(2)	Metoclopramide
	(3)	Diclofenac	(4)	Salicylic acid
67.	Which of the following drugs is used as osmotic diuretic?			
	(1)	Furosemide	(2)	Lasix
	(3)	Mannitol	(4)	Hydrochlorothiazide
68.	Mebendazole is used for treatment of the following infections except :			
	(1)	W. bancrofti	(2)	Hookworm
	(3)	Roundworm	(4)	Whipworm
69.	All of the following are rapid acting insulin $except$:			
	(1)	Insulin lispro	(2)	Regular insulin
	(3)	Insulin aspart	(4)	Insulin glulisne
70.	Honey and steam are natural remedies for relieve symptoms of :			
	(1)	Vomit	(2)	Fever
	(3)	Cough	(4)	Pain

BNS-042 (15)

Rough Work