

**POST BASIC BACHELOR OF SCIENCE
(NURSING) B.Sc. (N) (PB)**

Term-End Examination

June, 2014

BNS - 106 : MEDICAL SURGICAL NURSERY

Time : 3 hours

Maximum Marks : 70

Note : Attempt all Questions :

1. (a) Define chronic Renal failure and list its causes. 2+2+8
+3=15

(b) Write the nursing management of a patient with chronic renal failure

(c) Discuss the nursing responsibilities of a patient on haemodialysis

OR

(a) List four common causes of epilepsy 2+4+7
+2=15

(b) Explain the pathophysiology of epilepsy

(c) Describe the nursing management of a patient during a seizure

(d) List the steps for prevention of epilepsy

2. Briefly write the *nursing interventions* in *any three* of the following: **5x3=15**
- (a) A patient with C.S.F. formula
 - (b) A patient suffering from Ulcerative colitis
 - (c) A patient having acute pancreatitis
 - (d) A patient Undergone cardiac surgery
3. (a) Define cushing's syndrome **2+3+5+5=15**
- (b) List its causes and risk factors
 - (c) What clinical manifestations the patient is likely to have?
 - (d) Describe the nursing management for patient suffering from cushing's syndrome.
- 4 Write short notes on *any three* of the following: **5x3=15**
- (a) Pneumothorax
 - (b) Bronchial Asthma
 - (c) Respiratory Acidosis
 - (d) Counselling in HIV/AIDS
5. Fill in the blanks :
- (a) When the Ph rises above 7.45, the client has **1x5=5**
.....
 - (b) The diseases that occur as a result of immune responses against self antigens are termed as
.....

- (c) A sign of cardiopulmonary arrest, which appears later is
- (d) A period of isolation to prevent spread of infection is called as
- (e) The speech centre is situated in

6. Write 'T' if the statement is true and 'F' if it is false in your answer sheet.

- (a) IABP therapy reduce the work of left vascular by decreasing systemic vascular resistance. 1x5=5
- (b) Persons taking levodopa should also take vitamin B₆
- (c) Pressure on the median nerve leads to a condition called carpal Tunnel Syndrome
- (d) Cervical spinal cord injury results in Quadriplegia
- (e) Aphasia is usually seen in patients with right sided paralysis

— ** * —